

QUESTIONS?: PROCEDURE SCHEDULING: 352-331-8902 x 206

# Bravo pH Capsule/Upper Endoscopy Prep Instructions

Use this guide to help you prepare for your procedure appointment. Read the entire guide before beginning.

Physician.

110000010 20001	_ 11/51010111
Arrival Time:	Procedure Time:
Procedure scheduled at:	
Endoscopy Center at North Florid 6500 W. Newberry Road, 3 <sup>rd</sup> Floo You must call to pre-register 888-821-1632	
week before your appointment. If you haven't rea	our procedure. Some steps in this guide begin up to one ad through the guide at least one week in advance, call your mportaint. If you have questions, remember that we're here
If you don't follow all the steps listed he	ere, your procedure may be cancelled.
What does this guide cover?	
The day of your procedureWhat to expect, procedure charges and car	page 2 page 3 ncellation/missed appointmentpage 4 doscopypage 5

Procedure Date:

# 5 days before your procedure

Do I continue to take my Proton Pump Inhibitor (Nexium, Aciphex, Prevacid or Protonix)? STOP CONTINUE

If you have been instructed to stop taking your proton pump inhibitor, do so for 5 days prior to your procedure.

Blood Thinners such as Plavix, warfarin (Coumadin), or Lovenox

We will contact your doctor who prescribed the blood thinner medication to get approval to temporarily stop your medication for days before your procedure.		
Your labs will be drawn the day before your procedure to check you levels. After your procedure, your doctor will inform you when to restart your Coumadin.		
STOP TAKING: Aspirin products (Buffererin, Anacin, Ecotrin, Excedrin, Alka Seltzer, Goody's Powders, Elmiron, Pletal or Aggrenox		
Fish Oil Iron		
You may continue to take  ☑ All other medications as directed ☑ Acetaminophen (Tylenol) ☐ Other:		
Follow special instructions if you have any of these conditions:		
Follow special instructions if you have any of these conditions:  Diabetes Patients  Continue medication as prescribed		
<u>Diabetes Patients</u>		
Diabetes Patients Continue medication as prescribed  Implanted cardio defibrillator patients		

352-331-8902 x 206.

## The Day of your procedure

All prep medication must be finished at least 8 hours before your scheduled procedure.

Diabetic patients: Do not take your morning dose of insulin or antidiabetic pills today. Bring ALL medication with you to your procedure.



If you take medication for your heart, stomach, seizure, depression, or nerves, take your normal dose with a sip of water at least 4 hours before your scheduled time.

Bring any prescribed respiratory inhaler with you.

## **NO FOOD AFTER MIDNIGHT**

## CLEAR LIQUIDS are allowed up to 4 hours prior to your procedure.

A liquid is considered clear if you can read a newspaper through the glass.

**☑** water

☑ plain tea & coffee

☑ Gatorade, PowerAde, G2

**JUICES** 

✓ white cranberry

**☑** apple

✓ white grape

☑ filtered limeade, lemonade

**SOUPS** 

dear broth, consume

**DESSERT** 

**☑** gelatin

✓ water ices, popsicles

☑ clear sorbets (non-dairy)

NOTICE: Because the capsule contains a small magnet, patients should not have an MRI study within 30 days of undergoing the Bravo™ pH test

# **Checklist**

Use this checklist to help you remember everything you need to bring with you today.

- A friend or relative over 18 years old to drive you home (you cannot take a taxi or public transit home; you will be woozy from medication). The responsible party must remain on hospital grounds. You must not drive for 16 hours after the procedure.
- ☐ Complete Yellow Anesthesia Questionnaire Form. Bring completed form to the hospital the day of your procedure.
- ☐ Photo identification (license or other form of ID)
- Insurance cards
- ☐ Funds for deductible or co-payments (credit card, check etc.)
- ☐ Remove ALL piercings
- ☐ All medications in Rx bottle with label
- ☐ Other

## **PROCEDURE CHARGES:**

In addition to the facility fee, you will receive a bill from Digestive Disease Associates (a section of SIMEDHealth) – this will include:

- Anesthesia Fee
- Procedure Fee
- Pathology Fee (if biopsies are taken)

All insurance questions should be directed to our billing department (352-331-8902 X 202).

### **CANCELLATION OR MISSED APPOINTMENTS:**

Should you have to cancel and reschedule your procedure, a 48-hour (minimum) advance notice is required. Those that cancel with less than a 48-hour notice will be charged a \$100 missed procedure fee.



### INFORMATION REGARDING GASTROINTESTINAL ENDOSCOPY

### **BRIEF DESCRIPTION OF PROCEDURES:**

Gastrointestinal Endoscopy is the examination of the digestive tract with lighted instruments. At the time of the examination, the inside lining of the G.I. tract will be inspected thoroughly and may be photographed. A small portion of tissue may be removed for microscopic study (biopsy), or the tissue may be brushed or washed to collect cells for a special study. Polyps may be removed. A narrowed portion of the digestive tract can be stretched or dilated to a more normal size

**EGD** (Esophagogastroduodenoscopy) is the examination of the esophagus, stomach and duodenum.

**ESOPHAGEAL DILATION** is the stretching of a narrowed portion of the esophagus with a dilator.

FLEXIBLE SIGMOIDOSCOPY is the examination of the anus, rectum, and left lower colon.

**ENTEROSCOPY** is the examination of the small intestine.

**COLONOSCOPY** is the examination of the entire colon (large intestine)

POLYPECTOMY is the removal of small growths called polyps, with the use of either a wire loop and electric current or a cold forceps.

**BANDING** is the application of tiny bands to an area to reduce the risk of bleeding.

INJECTION THERAPY is the injection of a medication or solution to treat or mark an area.

**BRAVO** is the temporary placement of a capsule in the lower esophagus to record pH acid levels. The capsule will pass out of the G.I. tract naturally within 48 hours.

ANESTHESIA is the medication administered to achieve a level of moderate to deep sedation as deemed necessary by a nurse anesthetist and physician.

**THE PROCEDURE** — The endoscopy will be performed with you lying on your left side. Medications will be administered through the intravenous line and oxygen will be delivered into your nose. Blood pressure, breathing, and heart rhythm will be monitored. During upper endoscopy a plastic mouth guard will be placed between your teeth to prevent damage to your teeth and the scope. During colonoscopy a rectal exam will be performed.

**ALTERNATIVES**: Possible alternatives vary from each patient, as some alternatives may be inappropriate for a number of reasons. However, these potential alternatives include x-rays CT (virtual colonoscopy), surgery, no examination, stool tests for blood, Cologuard for screening, or other alternatives that have been explained to you.

POSSIBLE RISK AND COMPLICATIONS: These vary in frequency among different procedures but may include:

1. Infection

8. Dental damage

2. Bleeding

9. Irritation in vein at site of the IV line

Irregular heart beat

10. Death

- 4. Pneumonia
- 5. Rare possibility of a ruptured spleen
- 6. Injury to the lining of the intestinal tract may result in a hole (perforation) of the wall
- 7. Reaction to medication used for anesthesia, including cardiopulmonary arrest (stopping of heartbeat or breathing)

You should understand that the doctor cannot tell you about every possible risk, alternative, complication or side effect, but we did discuss the major ones. The practice of medicine is not an exact science. No guarantees have been made to you concerning the results of the procedure. The miss rate of colonoscopy ranges in various studies between 5% - 10%. Quality and accuracy of the exam is greatly enhanced by proper bowel cleansing. If bowel preparation is suboptimal you may be advised to return for a repeat exam within a short time interval.

**RECOVERY** — After the endoscopy, you will be kept for a time for observation while some of the medicine wears off. The most common discomfort after the examination is a feeling of bloating from the air introduced during the examination, which resolves quickly. You may be allowed to drink liquids during recovery. The medicines leave many patients feeling tired afterwards or they may have difficulty concentrating, so you cannot return to work that day.

The endoscopist can usually tell you the results of your examination before you leave the endoscopy unit. If biopsies have been taken or polyps removed, you will be instructed to call back for results. Tissue that has been removed is sent to a laboratory for analysis and it may take several days for a report to be completed.

**AFTER ENDOSCOPY** — Though patients worry about the discomforts of the examination, most people tolerate it very well and feel fine afterwards. Some fatigue is common after the examination, and you should plan to take it easy and relax the rest of the day.

You should contact your doctor about the results of your test if you have any questions and especially if biopsies were taken. The endoscopy team can give you some guidelines as to when your doctor should have all the results and whether further treatment will be necessary.

The following symptoms should be reported immediately: Severe abdominal pain (not just gas cramps), A firm, distended abdomen, Vomiting or Fever, Difficulty swallowing/severe sore throat, A crunching feeling under the skin, Rectal bleeding more than a few tablespoons

My signature below indicates that I have read and understand this handout. I have had the opportunity to ask questions about the procedure and my questions, if any, have been answered to my satisfaction.

My signature below also indicates that if I have a biopsy taken during my procedure, I should hear from the office in about 14 day if not I will call in 14 days (\_\_\_\_\_\_\_) to get my results.

I have received a North Florida Endoscopy Center packet, which includes: ◆ Patient rights and responsibilities, ◆ Disclosure of physician ownership in the Endoscopy Center, ◆ Information and facility policies concerning advanced directives, ◆ Information concerning the patient informed consent.



### INFORMATION REGARDING GASTROINTESTINAL ENDOSCOPY

### **BRIEF DESCRIPTION OF PROCEDURES:**

Gastrointestinal Endoscopy is the examination of the digestive tract with lighted instruments. At the time of the examination, the inside lining of the G.I. tract will be inspected thoroughly and may be photographed. A small portion of tissue may be removed for microscopic study (biopsy), or the tissue may be brushed or washed to collect cells for a special study. Polyps may be removed. A narrowed portion of the digestive tract can be stretched or dilated to a more normal size

**EGD** (Esophagogastroduodenoscopy) is the examination of the esophagus, stomach and duodenum.

**ESOPHAGEAL DILATION** is the stretching of a narrowed portion of the esophagus with a dilator.

FLEXIBLE SIGMOIDOSCOPY is the examination of the anus, rectum, and left lower colon.

**ENTEROSCOPY** is the examination of the small intestine.

**COLONOSCOPY** is the examination of the entire colon (large intestine)

POLYPECTOMY is the removal of small growths called polyps, with the use of either a wire loop and electric current or a cold forceps.

**BANDING** is the application of tiny bands to an area to reduce the risk of bleeding.

INJECTION THERAPY is the injection of a medication or solution to treat or mark an area.

BRAVO is the temporary placement of a capsule in the lower esophagus to record pH acid levels. The capsule will pass out of the G.I. tract naturally

ANESTHESIA is the medication administered to achieve a level of moderate to deep sedation as deemed necessary by a nurse anesthetist and physician.

THE PROCEDURE — The endoscopy will be performed with you lying on your left side. Medications will be administered through the intravenous line and oxygen will be delivered into your nose. Blood pressure, breathing, and heart rhythm will be monitored. During upper endoscopy a plastic mouth guard will be placed between your teeth to prevent damage to your teeth and the scope. During colonoscopy a rectal exam will be performed.

ALTERNATIVES: Possible alternatives vary from each patient, as some alternatives may be inappropriate for a number of reasons. However, these potential alternatives include x-rays CT (virtual colonoscopy), surgery, no examination, stool tests for blood, Cologuard for screening, or other alternatives that have been explained to you.

Irritation in vein at site of the IV line

8. Dental damage

POSSIBLE RISK AND COMPLICATIONS: These vary in frequency among different procedures but may include:

- 1. Infection
- 2.
  - Bleeding 9.
- Irregular heart beat
- 4. Pneumonia
- Rare possibility of a ruptured spleen
- Injury to the lining of the intestinal tract may result in a hole (perforation) of the wall 6.
- Reaction to medication used for anesthesia, including cardiopulmonary arrest (stopping of heartbeat or breathing)

10. Death

You should understand that the doctor cannot tell you about every possible risk, alternative, complication or side effect, but we did discuss the major ones. The practice of medicine is not an exact science. No guarantees have been made to you concerning the results of the procedure. The miss rate of colonoscopy ranges in various studies between 5% - 10%. Quality and accuracy of the exam is greatly enhanced by proper bowel cleansing. If bowel preparation is suboptimal you may be advised to return for a repeat exam within a short time interval.

RECOVERY — After the endoscopy, you will be kept for a time for observation while some of the medicine wears off. The most common discomfort after the examination is a feeling of bloating from the air introduced during the examination, which resolves quickly. You may be allowed to drink liquids during recovery. The medicines leave many patients feeling tired afterwards or they may have difficulty concentrating, so you cannot return to work that day.

The endoscopist can usually tell you the results of your examination before you leave the endoscopy unit. If biopsies have been taken or polyps removed, you will be instructed to call back for results. Tissue that has been removed is sent to a laboratory for analysis and it may take several days for a report to be completed.

AFTER ENDOSCOPY — Though patients worry about the discomforts of the examination, most people tolerate it very well and feel fine afterwards. Some fatigue is common after the examination, and you should plan to take it easy and relax the rest of the day.

You should contact your doctor about the results of your test if you have any questions and especially if biopsies were taken. The endoscopy team can give you some guidelines as to when your doctor should have all the results and whether further treatment will be necessary.

The following symptoms should be reported immediately: Severe abdominal pain (not just gas cramps), A firm, distended abdomen, Vomiting or Fever, Difficulty swallowing/severe sore throat, A crunching feeling under the skin, Rectal bleeding more than a few tablespoons

My signature below indicates that I have read and understand this handout. I have had the opportunity to ask questions about the procedure and my questions, if any, have been answered to my satisfaction. My signature below also indicates that if I have a biopsy taken during my procedure, I should hear from the office in about 14 day if not I will call in 14 days (\_ \_) to get my results. I have received a North Florida Endoscopy Center packet, which includes: • Patient rights and responsibilities, • Disclosure of physician ownership in the Endoscopy Center, ◆ Information and facility policies concerning advanced directives, ◆ Information concerning the patient informed consent.

	<u>X</u>
Print Patient's Name	Patient's Signature
Witness' Signature	Date
DDA Account Number	Procedure Date

Page | 7 Bravo pH Capsule